









7 Clinical Stages of Alzheimer's

Alzheimer's disease is the most common cause of dementia, a neurological brain disorder that causes problems with memory.




The Seven Clinical Stages of Dementia, also known as the Global Deterioration Scale (GDS), was developed by Dr. Barry Reisberg, Director of the Fisher Alzheimer's Disease Education and Research program at NYU Grossman School of Medicine. This guideline is used




by professionals and caregivers around the world to identify what stage of the disease a person is in. Stages 1-3 are the pre-dementia stages; stages 4-7 are the dementia stages. Stage 5 is the point where a person can no longer live without assistance.

STAGE 1 NO DEMENTIA SEEN




STAGE 2 SUBJECTIVE MEMORY LOSS AGE RELATED FORGETFULNESS



STAGE 3 MILD COGNITIVE IMPAIRMENT

The person begins to show subtle changes in concentration, thinking, comprehension, or memory-which may be noticeable by family and friends. They may begin to repeat statements, and there is a decline in their work performance if they are still working.



STAGE 4 MODERATE COGNITIVE DECLINE MILD DEMENTIA

The person's cognitive deficits are more pronounced, and several of their functions are affected. They have poor short-term memory, and frequently have difficulty with managing their personal finances.



STAGE 5

MODERATELY SEVERE COGNITIVE DECLINE MODERATE DEMENTIA

The person may no longer have the ability to live independently. They may exhibit the inability to recall very familiar information such as their phone number of many years, close relatives or close friend's names, and may have difficulty selecting proper clothing for the season.



STAGE 6 SEVERE COGNITIVE DECLINE MODERATELY SEVERE DEMENTIA





STAGE 7 VERY SEVERE COGNITIVE DECLINE SEVERE DEMENTIA

The person is in the final stage of dementia and will need constant care with all daily activities. Early in this stage they may be able to speak a half-dozen intelligible words. As this stage progresses, physical motor skills are compromised to the point where the person can no longer walk independently. Later they lose the ability to sit up and to smile.

This disease can affect anyone. It is imperative that when a diagnosis is given, efforts are made to obtain as much information as possible for care and support. With proper care, the person with dementia can remain content.



For more detailed information about the 7 Clinical Stages of Alzheimer's go to: alzinfo.org/understand-alzheimers



*Approximate years are based on healthy persons



someone in the U.S. develops
Alzheimer's disease

In the U.S. alone, more than 11 million patients receive care from unpaid loved ones who help with feeding, dressing, bathing, administering medications, and other activities of daily living.

MORE THAN 6 MILLION AMERICANS ARE LIVING WITH ALZHEIMER'S TODAY



FDR Station, PO Box 220 New York, NY 10150

1.800.ALZ.INFO (1.800.259.4636) 1.212.915.1328 Email: info@alzinfo.org

Website: alzinfo.org



Together <u>We Can End Alzhei</u>mer's

FOLLOW US ON



/alzresearch



/fishercenter



/alzinfo



/company/fishercenter



We are the proud recipient of Charity Navigator's highest 4-Star ranking and Guide Star's Gold Seal of Transparency

